arXiv:1401.5498v1 [math.PR] 21 Jan 2014

Excursion Probability of Gaussian Random Fields on Sphere

Dan Cheng
North Carolina State University

Yimin Xiao *
Michigan State University

January 23, 2014

Abstract

Let $X = \{X(x) : x \in \mathbb{S}^N\}$ be a real-valued, centered Gaussian random field indexed on the N-dimensional unit sphere $\mathbb{S}^N$. Approximations to the excursion probability $\mathbb{P}\big\{\sup_{x\in\mathbb{S}^N} X(x) \geq u\big\}$, as $u \to \infty$, are obtained for two cases: (i) X is locally isotropic and its sample path is non-smooth and; (ii) X is isotropic and its sample path is twice differentiable. For case (i), it is shown that the asymptotics is similar to Pickands' approximation on the Euclidean space which involves Pickands' constant; while for case (ii), we use the expected Euler characteristic method to obtain a more precise approximation such that the error is super-exponentially small.

Key Words: Excursion probability, Gaussian random fields on sphere, Pickands' constant, Euler characteristic.

1 Introduction

Even though the characterizations of isotropic covariance functions and variograms on spheres were given long time ago by Schoenberg (1942) and Gangolli (1967), respectively, and random fields on the sphere were studied by Obukhov (1947), Yaglom (1961) and Jones (1963), it is the applications in atmospherical sciences, geophysics, solar physics, medical imaging and environmental sciences [see, e.g., Genovese et al. (2004), Oh and Li (2004), Stein (2007), Cabella and Marinucci (2009), Tebaldi and Sansó (2009), Hansen et al. (2012)] that have stimulated the recent rapid developments in statistics of random fields on the sphere. Various new random field models have been constructed and new probabilistic and statistical methods have been developed. For example, Jun and Stein (2007, 2008), Huang et al. (2009), Jun (2011), Hitczenko and Stein (2012), Ma (2012), Du et al. (2013) and Gneiting (2013) have constructed several classes of real or vector-valued random fields on spheres; Istas (2005,

*Research partially supported by NSF grants DMS-1309856 and DMS-1307470.

2006) has constructed spherical fractional Brownian motion (SFBM), which has fractal sample functions, and studied its Karhunen-Loève expansion and other properties. Lang and Schwab (2013) characterized sample Hölder continuity and sample differentiability of isotropic Gaussian random fields on the two-dimensional sphere $\mathbb{S}^2$ in terms of their angular power spectra. We refer to the recent book by Marinucci and Peccati (2011) for a systematic account on theory and statistical inferences of random fields on the sphere $\mathbb{S}^N$, with a view towards applications to cosmology.

In this paper we consider a real-valued, centered isotropic Gaussian random field $X = \{X(x) : x \in \mathbb{S}^N\}$, indexed on the N-dimensional unit sphere $\mathbb{S}^N$, and investigate the asymptotic properties of the excursion probability $\mathbb{P}\big\{ \sup_{x\in\mathbb{S}^N} X(x) \geq u\big\}$ as $u \to \infty$. Such excursion probabilities are important in probability theory, statistics and their applications. In particular, we mention that the above excursion probability has appeared in Sun (1991), Park and Sun (1998) to determine the P-value for studying exploratory projection pursuit and, as illustrated by Sun (2001), is useful for constructing simultaneous confidence region for a function $f : \mathbb{S}^N \to \mathbb{R}$. In his studies of projection-based depth functions, Zuo (2003) has shown that Gaussian random fields on sphere appear as scaling limit of sample projection median [see Theorems 3.2 and 3.3 in Zuo (2003)] and the excursion probability of the limiting Gaussian field is useful for constructing confidence regions for the true projection median [see Remark 3.2 in Zuo (2003)]. For further information on extreme value theory of Gaussian random fields on Euclidean spaces or manifolds and statistical applications, we refer to Adler and Taylor (2007) and Adler et al. (2012).

We will distinguish two cases: (i) the sample path of X is non-smooth and, (ii) $X(\cdot) \in C^2$ a.s. For the non-smooth case, our argument is based on the extension of the asymptotic results of Pickands (1969) and Piterbarg (1996) by Chan and Lai (2006). The main result for Case (i) is Theorem 2.4, where one may observe that the asymptotics is very similar to the classical Pickands' approximation for locally isotropic Gaussian fields on the Euclidean space [see Theorem 2.2]. The method for deriving Theorem 2.4 can also be applied to other Gaussian fields on sphere with more complicated local covariance structures, see Section 2.2 for the example of standardized spherical fractional Brownian motion. For the smooth case, we consider isotropic Gaussian fields on sphere. Thanks to the special representation of covariance function [Theorem 3.1], we are able to apply the general theory of Adler and Taylor (2007) to compute the Lipschitz-Killing curvatures induced by the field and hence derive the approximation to the excursion probability, see Theorem 3.7 and Corollary 3.9 below. Such approximation is more precise than that in Theorem 2.4 for the non-smooth case and the error is super-exponentially small.

We should mention that Mikhaleva and Piterbarg (1997) have established asymptotic results for the excursion probability of Gaussian random fields on a finite dimensional smooth

manifold in $\mathbb{R}^{N+1}$. Their theorems can be applied to obtain results similar to Theorem 2.4 in this paper for a Gaussian random field X on the sphere $\mathbb{S}^N$, provided X is the restriction on $\mathbb{S}^N$ of a Gaussian random field defined on $\mathbb{R}^{N+1}$ and satisfies the conditions on the covariance structures (i.e., the local stationarity condition) in Mikhaleva and Piterbarg (1997). Since not every Gaussian random field on $\mathbb{S}^N$ can be obtained as the restriction of a Gaussian random field on $\mathbb{R}^{N+1}$ [cf. e.g., Ma (2012, p. 775)] and the conditions on the covariance structure in Mikhaleva and Piterbarg (1997) are not always easy to verify, we have decided in the present paper to deal with Gaussian random fields on sphere directly.

For $x = (x_1, \ldots, x_{N+1}) \in \mathbb{S}^N$, its corresponding spherical coordinate $\theta = (\theta_1, \ldots, \theta_N)$ is defined as follows.

$$\begin{aligned} x_1 &= \cos\theta_1, \\ x_2 &= \sin\theta_1 \cos\theta_2, \\ x_3 &= \sin\theta_1 \sin\theta_2 \cos\theta_3, \\ &\vdots \\ x_N &= \sin\theta_1 \sin\theta_2 \cdots \sin\theta_{N-1} \cos\theta_N, \\ x_{N+1} &= \sin\theta_1 \sin\theta_2 \cdots \sin\theta_{N-1} \sin\theta_N, \end{aligned} \tag{1.1}$$

where $0 \le \theta_i \le \pi$ for $1 \le i \le N-1$ and $0 \le \theta_N < 2\pi$.

Throughout this paper, for two points $x = (x_1, \ldots, x_{N+1})$ and $y = (y_1, \ldots, y_{N+1})$ on $\mathbb{S}^N$, we always denote by $\theta = (\theta_1, \ldots, \theta_N)$ the spherical coordinate of x and by $\varphi = (\varphi_1, \ldots, \varphi_N)$ the spherical coordinate of y respectively.

Let $\|\cdot\|$, $\langle \cdot, \cdot \rangle$ be respectively the Euclidean norm and the inner product in $\mathbb{R}^{N+1}$ (or in $\mathbb{R}^N$, which will be clear from the context). Denote by $d(\cdot, \cdot)$ the spherical distance on $\mathbb{S}^N$, i.e., $d(x, y) = \arccos\langle x, y \rangle$, $\forall x, y \in \mathbb{S}^N$. For two functions $f(t)$ and $g(t)$, we say $f(t) \sim g(t)$ as $t \to t_0 \in [-\infty, +\infty]$ if $\lim_{t \to t_0} f(t)/g(t) = 1$.

2 Non-smooth Gaussian Fields on Sphere

2.1 Locally Isotropic Gaussian Fields on Sphere

Let $X = \{X(x) : x \in \mathbb{S}^N\}$ be a centered Gaussian random field with covariance function C satisfying

$$C(x, y) = 1 - c d^{\alpha}(x, y)(1 + o(1)) \quad \text{as } d(x, y) \to 0, \tag{2.1}$$

for some constants $c > 0$ and $\alpha \in (0, 2]$.

Covariance functions satisfying (2.1) behave isotropically in a local sense, hence the corresponding random fields fall under the general category of locally isotropic random fields.

Similarly to Gaussian random fields defined on the Euclidean space [cf. Adler (2010)], one can show that, when $\alpha \in (0,2)$, the sample function of X is not differentiable and the fractal dimensions of its trajectories are determined by α. See Andreev and Lang (2013), Hansan et al. (2012) and Lang and Schwab (2013) for related results.

There are many examples of covariances of isotropic Gaussian fields on $\mathbb{S}^N$ satisfying (2.1). A well-known example is $C(x,y) = e^{-cd^\alpha(x,y)}$, where $c > 0$ and $\alpha \in (0,1]$ [cf. e.g., Huang et al. (2011, p. 725)]. In their studies on germ-grain (or random balls) models on the sphere $\mathbb{S}^N$, Estrade and Istas (2010, Remark 2.5 and Lemma 3.1) discovered an isotropic Gaussian random field W^β on $\mathbb{S}^N$ with $0 < \beta < 1/2$, whose covariance function satisfies (2.1) for $\alpha = 2\beta \in (0,1]$. [From here one can show that, even though W^β and the spherical fractional Brownian motion $B_\beta(x)$ introduced by Istas (2005) are different, they share some local properties (e.g., they have the same Hölder continuity and fractal dimensions). In Remark 2.6 below, we will compare the excursion probabilities of W^β and the standardized SFBM.] Moreover, as in Yadrenko (1983) and Ma (2012), one can apply the identity

$$\|x - y\| = 2\sin\Big(\frac{d(x,y)}{2}\Big), \quad \forall x,\, y \in \mathbb{S}^N$$

to construct covariance functions that satisfy (2.1) from isotropic covariance functions $K(\cdot)$ on $\mathbb{R}^N$ which satisfy $K(x) = 1 - c_1\|x\|^\alpha(1 + o(1))$ as $\|x\| \to 0$. In particular, the following covariance function C given by Soubeyrand et al. (2008)

$$C(x,y) = 1 - \left(\sin\frac{d(x,y)}{c^{1/\alpha}}\right)^\alpha \mathbb{1}_{\{d(x,y)\le \pi c^{1/\alpha}\}}, \tag{2.2}$$

where $c > 0$ and $\alpha \in (0,2)$ are constants, satisfies (2.1). See Huang et al. (2011) and Gneiting (2013) for further comments on (2.2) and more examples.

Recall the spherical coordinate representation in (1.1), we define the Gaussian random field $\widetilde{X} = \{\widetilde{X}(\theta) : \theta \in [0,\pi]^{N-1} \times [0,2\pi)\}$ by $\widetilde{X}(\theta) := X(x)$ and denote by $\widetilde{C}$ the covariance function of $\widetilde{X}$ accordingly.

The following lemma characterizes the local behavior of spherical distance of two points on sphere. It provides a useful tool to establish the relation between the local behavior of the covariance functions C and $\widetilde{C}$.

Lemma 2.1 *Let $x, y \in \mathbb{S}^N$ and let x be fixed. Then as $d(y,x) \to 0$,*

$$d^2(y,x) \sim (\varphi_1 - \theta_1)^2 + (\sin^2\theta_1)(\varphi_2 - \theta_2)^2 + \cdots + \Big(\prod_{i=1}^{N-1}\sin^2\theta_i\Big)(\varphi_N - \theta_N)^2,$$

where $\theta = (\theta_1, \ldots, \theta_N)$ and $\varphi = (\varphi_1, \ldots, \varphi_N)$ are the spherical coordinates of x and y respectively.

Proof For $x, y \in \mathbb{S}^N$, we see that $d(y,x) \to 0$ implies $\|y - x\| \to 0$ and

$$\cos \|y - x\| \sim 1 - \frac{1}{2}\|y - x\|^2 = \langle y, x \rangle.$$

Then as $d(y,x) \to 0$,

$$\begin{aligned}
d^2(y,x) &= \arccos^2 \langle y, x \rangle \sim \|y - x\|^2 \\
&= (\cos\varphi_1 - \cos\theta_1)^2 + (\sin\varphi_1 \cos\varphi_2 - \sin\theta_1 \cos\theta_2)^2 + \cdots \\
&\quad + (\sin\varphi_1 \sin\varphi_2 \cdots \sin\varphi_{N-1} \cos\varphi_N - \sin\theta_1 \sin\theta_2 \cdots \sin\theta_{N-1} \cos\theta_N)^2 \\
&\quad + (\sin\varphi_1 \sin\varphi_2 \cdots \sin\varphi_{N-1} \sin\varphi_N - \sin\theta_1 \sin\theta_2 \cdots \sin\theta_{N-1} \sin\theta_N)^2 \\
&= 2 - 2\cos(\varphi_1 - \theta_1) + 2(\sin\varphi_1 \sin\theta_1)[1 - \cos(\varphi_2 - \theta_2)] \\
&\quad + \cdots + 2\Big(\prod_{i=1}^{N-1} \sin\varphi_i \sin\theta_i\Big)[1 - \cos(\varphi_N - \theta_N)].
\end{aligned}$$

It follows from the spherical coordinates that $d(y,x) \to 0$ is equivalent to $\|\varphi - \theta\| \to 0$. (There is an exception for θ with $\theta_N = 0$, since for those φ such that $d(y,x) \to 0$ and φ_N tending to 2π, $\|\varphi - \theta\|$ does not tend to 0. In such case, we may treat θ_N as 2π instead of 0 and this does not affect the result thanks to the periodicity.) Therefore, as $d(y,x) \to 0$, by Taylor's expansion,

$$\begin{aligned}
d^2(y,x) &\sim (\varphi_1 - \theta_1)^2 + (\sin\varphi_1 \sin\theta_1)(\varphi_2 - \theta_2)^2 + \cdots + \Big(\prod_{i=1}^{N-1} \sin\varphi_i \sin\theta_i\Big)(\varphi_N - \theta_N)^2 \\
&\sim (\varphi_1 - \theta_1)^2 + (\sin^2\theta_1)(\varphi_2 - \theta_2)^2 + \cdots + \Big(\prod_{i=1}^{N-1} \sin^2\theta_i\Big)(\varphi_N - \theta_N)^2.
\end{aligned}$$

This completes the proof. □

Next, we recall from Chan and Lai (2006) some results on approximations to the excursion probability of Gaussian random fields over the Euclidean space.

Let $0 < \alpha \le 2$ and let $\{W_t(s) : s \in [0,\infty)^N\}$ $(t \in \mathbb{R}^N)$ be a family of Gaussian random fields such that

$$\begin{aligned}
\mathbb{E}W_t(s) &= -\|s\|^\alpha r_t(s/\|s\|), \\
\mathrm{Cov}\big(W_t(s), W_t(v)\big) &= \|s\|^\alpha r_t(s/\|s\|) + \|v\|^\alpha r_t(v/\|v\|) \\
&\qquad - \|s - v\|^\alpha r_t((s-v)/\|s-v\|),
\end{aligned} \tag{2.3}$$

where $r_t(\cdot) : \mathbb{S}^{N-1} \to \mathbb{R}_+$ is a continuous function which satisfies

$$\sup_{v \in \mathbb{S}^{N-1}} |r_t(v) - r_s(v)| \to 0 \quad \text{as } s \to t. \tag{2.4}$$

Define

$$H_\alpha^r(t) = \lim_{K\to\infty} K^{-N} \int_0^\infty e^u \mathbb{P}\Big\{ \sup_{s\in[0,K]^N} W_t(s) \geq u \Big\} du. \tag{2.5}$$

Denote by H_α the usual Pickands' constant, that is

$$H_\alpha = \lim_{K\to\infty} K^{-N} \int_0^\infty e^u \mathbb{P}\Big\{ \sup_{s\in[0,K]^N} Z(s) \geq u \Big\} du,$$

where $\{Z(s) : s \in [0,\infty)^N\}$ is a Gaussian random field such that

$$\mathbb{E}Z(s) = -\|s\|^\alpha,$$
$$\mathrm{Cov}(Z(s), Z(v)) = \|s\|^\alpha + \|v\|^\alpha - \|s-v\|^\alpha.$$

It is clear that $H_\alpha^r(t)$ becomes H_α when $r_t \equiv 1$.

Let $D \subset \mathbb{R}^N$ be a bounded N-dimensional Jordan measurable set, that is, the boundary of D has N-dimensional Lebesgue measure 0. Let $Y = \{Y(t), t \in \mathbb{R}^N\}$ be a real-valued, centered Gaussian field such that its covariance function C_Y satisfies

$$C_Y(t, t+s) = 1 - \|s\|^\alpha r_t(s/\|s\|)(1+o(1)) \quad \text{as } \|s\| \to 0, \tag{2.6}$$

for some constant $\alpha \in (0, 2]$, uniformly over $t \in \bar{D}$, the closure of D.

We will make use of the following theorem of Chan and Lai (2006). See also Piterbarg (1996), Mikhaleva and Piterbarg (1997) for similar results under somewhat more restrictive conditions.

Theorem 2.2 [Chan and Lai (2006, Theorem 2.1)] *Let $D \subset \mathbb{R}^N$ be a bounded N-dimensional Jordan measurable set. Suppose the Gaussian random field $\{Y(t) : t \in \mathbb{R}^N\}$ satisfies condition (2.6), in which $r_t(\cdot) : \mathbb{S}^{N-1} \to \mathbb{R}_+$ is a continuous function such that the convergence (2.4) is uniformly in $\bar{D}$ and $\sup_{t\in\bar{D}, v\in\mathbb{S}^{N-1}} r_t(v) < \infty$. Then as $u \to \infty$,*

$$\mathbb{P}\Big\{ \sup_{t\in D} Y(t) \geq u \Big\} \sim u^{2N/\alpha} \Psi(u) \int_D H_\alpha^r(t) dt.$$

Here and in the sequel, $\Psi(u) = (\sqrt{2\pi}\, u)^{-1} e^{-u^2/2}$.

The lemma below establishes the relation between $H_\alpha^r(t)$ and H_α for a special class of functions r.

Lemma 2.3 *Let $\{W_t(s) : s \in [0,\infty)^N\}$ ($t \in \mathbb{R}^N$) be a family of Gaussian random fields satisfying (2.3) with*

$$r_t(v) = \|M_t v\|^\alpha, \quad \forall v \in \mathbb{S}^{N-1},$$

where, for every $t \in \mathbb{R}^N$, M_t is a non-degenerate $N \times N$ matrix. Then for each $t \in \mathbb{R}^N$,

$$H_\alpha^r(t) = |\det M_t| H_\alpha.$$

Proof Let $t \in \mathbb{R}^N$ be fixed and consider the centered Gaussian random field $\overline{W}_t = \{\overline{W}_t(s),\, s \in [0,\infty)^N\}$ defined by $\overline{W}_t(s) = W_t(M_t^{-1}s)$. Then by (2.3), $\overline{W}_t$ satisfies

$$\begin{aligned} \mathbb{E}\overline{W}_t(s) &= -\|s\|^\alpha, \\ \mathrm{Cov}(\overline{W}_t(s), \overline{W}_t(v)) &= \|s\|^\alpha + \|v\|^\alpha - \|s-v\|^\alpha. \end{aligned} \tag{2.7}$$

Let $B_K = [0,K]^N$ and define $M_tB_K = \{s \in \mathbb{R}^N : \exists v \in B_K \text{ such that } s = M_tv\}$. Note that $\mathrm{Vol}(M_tB_K) = |\det M_t|\mathrm{Vol}(B_K)$ and $\sup_{s\in B_K} W_t(s) = \sup_{s\in M_tB_K} \overline{W}_t(s)$, it follows from (2.5) that

$$\begin{aligned} H_\alpha^r(t) &= \lim_{K\to\infty} \frac{1}{\mathrm{Vol}(B_K)} \int_0^\infty e^u \mathbb{P}\Big\{ \sup_{s\in B_K} W_t(s) \ge u \Big\} du \\ &= \lim_{K\to\infty} \frac{\mathrm{Vol}(M_tB_K)}{\mathrm{Vol}(B_K)} \frac{1}{\mathrm{Vol}(M_tB_K)} \int_0^\infty e^u \mathbb{P}\Big\{ \sup_{s\in M_tB_K} \overline{W}_t(s) \ge u \Big\} du \\ &= |\det M_t| \lim_{K\to\infty} \frac{1}{\mathrm{Vol}(M_tB_K)} \int_0^\infty e^u \mathbb{P}\Big\{ \sup_{s\in M_tB_K} \overline{W}_t(s) \ge u \Big\} du. \end{aligned}$$

Because of (2.7), we can modify the proofs in Qualls and Watanabe (1973) to show that

$$H_\alpha = \lim_{K\to\infty} \frac{1}{\mathrm{Vol}(M_tB_K)} \int_0^\infty e^u \mathbb{P}\Big\{ \sup_{s\in M_tB_K} \overline{W}_t(s) \ge u \Big\} du.$$

This completes the proof. □

For any $T \subset \mathbb{S}^N$, we denote by $D \subset [0,\pi]^{N-1} \times [0,2\pi)$ the set corresponding to T under the spherical coordinates (1.1). We say that T is an N-dimensional Jordan measurable set on $\mathbb{S}^N$ if D is an N-dimensional Jordan measurable set in $\mathbb{R}^N$. Now we can prove our main result of this section.

Theorem 2.4 *Let $\{X(x) : x \in \mathbb{S}^N\}$ be a centered Gaussian random field satisfying condition (2.1) and let $T \subset \mathbb{S}^N$ be an N-dimensional Jordan measurable set on $\mathbb{S}^N$. Then as $u \to \infty$,*

$$\mathbb{P}\Big\{ \sup_{x\in T} X(x) \ge u \Big\} \sim c^{N/\alpha} \mathrm{Area}(T) H_\alpha u^{2N/\alpha} \Psi(u),$$

where $\mathrm{Area}(T)$ *denotes the spherical area of T and $c > 0$ is the constant in (2.1).*

Proof For any $\theta \in [0,\pi]^{N-1} \times [0,2\pi)$, let M_θ be the $N \times N$ diagonal matrix $M_\theta = c^{1/\alpha}\mathrm{diag}(1, \sin\theta_1, \ldots, \prod_{i=1}^{N-1} \sin\theta_i)$. Here, we set $M_\theta = c^{1/\alpha}$ if $N = 1$. By Lemma 2.1, condition (2.1) implies

$$\widetilde{C}(\theta, \theta+\xi) = 1 - \|\xi\|^\alpha r_\theta(\xi/\|\xi\|)(1+o(1)) \quad \text{as } \|\xi\| \to 0,$$

where $r_\theta(\tau) = \|M_\theta\tau\|^\alpha$, $\forall \tau \in \mathbb{S}^{N-1}$. Then by Theorem 2.2, as $u \to \infty$,

$$\mathbb{P}\Big\{ \sup_{x\in T} X(x) \ge u \Big\} = \mathbb{P}\Big\{ \sup_{\theta\in D} \widetilde{X}(\theta) \ge u \Big\} \sim u^{2N/\alpha} \Psi(u) \int_D H_\alpha^r(\theta) d\theta. \tag{2.8}$$

It follows from Lemma 2.3 that for any $\theta \in [0,\pi]^{N-1}\times[0,2\pi)$ such that M_θ is non-degenerate (i.e., $\prod_{i=1}^{N-1}\sin\theta_i \neq 0$),

$$H_\alpha^r(\theta) = c^{N/\alpha}\bigg(\prod_{i=1}^{N-1}\sin^{N-i}\theta_i\bigg)H_\alpha.$$

Note that $(\prod_{i=1}^{N-1}\sin^{N-i}\theta_i)d\theta$ is the spherical area element and M_θ are non-degenerate for $\theta \in D$ almost everywhere, we obtain

$$\int_D H_\alpha^r(\theta)d\theta = c^{N/\alpha}\mathrm{Area}(T)H_\alpha.$$

Plugging this into (2.8) gives the desired result. □

Remark 2.5 Motivated by Mikhaleva and Piterbarg (1997), we think it would be interesting to study the excursion probability for Gaussian fields over Riemannian manifolds (beyond sphere) whose covariance functions satisfy (2.6) with $d(x,y)$ being the geodesic distance of x and y. We conjecture that Pickands-type approximation similar to Theorem 2.4 still holds.

2.2 Standardized Spherical Fractional Brownian Motion

Theorem 2.4 provides a nice approximation to the excursion probability for locally isotropic Gaussian random fields on $\mathbb{S}^N$ whose covariance functions satisfy (2.1). When the local behavior of the covariance function becomes more complicated, Theorem 2.4 may not be applicable anymore. However, we can still apply Lemma 2.1 to find the corresponding local behavior of covariance function under spherical coordinates and then apply Theorem 2.2 to obtain the asymptotics for the excursion probability. In the following, we use spherical fractional Brownian motion on sphere as an illustrating example.

Let o be a fixed point on $\mathbb{S}^N$. The spherical fractional Brownian motion (SFBM) $B_\beta = \{B_\beta(x) : x \in \mathbb{S}^N\}$ is defined by Istas (2005) as a centered real-valued Gaussian random field such that

$$B_\beta(o) = 0$$
$$\mathbb{E}(B_\beta(x) - B_\beta(y))^2 = d^{2\beta}(x,y), \quad \forall x,y \in \mathbb{S}^N,$$

where $\beta \in (0,1/2]$. It follows immediately that

$$\mathrm{Cov}(B_\beta(x), B_\beta(y)) = \frac{1}{2}\big(d^{2\beta}(x,o) + d^{2\beta}(y,o) - d^{2\beta}(x,y)\big).$$

Without loss of generality, we take $o = (1,0,\ldots,0) \in \mathbb{R}^{N+1}$, whose corresponding spherical coordinate is $(0,\ldots,0) \in \mathbb{R}^N$. We consider the standardized SFBM $X = \{X(x) : x \in \mathbb{S}^N\backslash\{o\}\}$ defined by

$$X(x) = \frac{B_\beta(x)}{d^\beta(x,o)}, \quad \forall x \in \mathbb{S}^N\backslash\{o\}. \tag{2.9}$$

Then the covariance of X is

$$C(x,y) = \mathrm{Cov}(X(x), X(y)) = \frac{d^{2\beta}(x,o) + d^{2\beta}(y,o) - d^{2\beta}(x,y)}{2d^{\beta}(x,o)d^{\beta}(y,o)}.$$

Note that under the spherical coordinates, $d(x,o) = \theta_1$ and $d(y,o) = \varphi_1$, together with Lemma 2.1, we obtain that the covariance function of corresponding Gaussian field $\widetilde{X}$ satisfies

$$\widetilde{C}(\theta,\varphi) = \mathrm{Cov}(\widetilde{X}(\theta), \widetilde{X}(\varphi)) = 1 - (1+o(1))\frac{1}{2\theta_1^{2\beta}}\Big[(\varphi_1-\theta_1)^2 + (\sin^2\theta_1)(\varphi_2-\theta_2)^2 + \cdots + \Big(\prod_{i=1}^{N-1}\sin^2\theta_i\Big)(\varphi_N-\theta_N)^2\Big]^{\beta}$$

as $d(x,y) \to 0$. Let

$$M_\theta = \frac{1}{2^{1/(2\beta)}\theta_1}\mathrm{diag}\Big(1, \sin\theta_1, \ldots, \prod_{i=1}^{N-1}\sin\theta_i\Big),$$
$$r_\theta(\tau) = \|M_\theta\tau\|^{2\beta}, \quad \forall \tau \in \mathbb{S}^{N-1},$$

and $\xi = \varphi - \theta$, then as $\|\xi\| \to 0$,

$$\widetilde{C}(\theta, \theta+\xi) = 1 - \|\xi\|^{2\beta} r_\theta(\xi/\|\xi\|)(1+o(1)).$$

Let $T \subset \mathbb{S}^N$ be an N-dimensional Jordan measurable set such that $o \notin \bar{T}$, and denote its corresponding domain under the spherical coordinates by D, which implies $\theta_1 \neq 0$ for any $\theta \in \bar{D}$. By Theorem 2.2, as $u \to \infty$,

$$\mathbb{P}\Big\{\sup_{x\in T} X(x) \geq u\Big\} = \mathbb{P}\Big\{\sup_{\theta\in D}\widetilde{X}(\theta) \geq u\Big\} \sim u^{N/\beta}\Psi(u)\int_D H_{2\beta}^r(\theta)d\theta.$$

It follows from Lemma 2.3 that for any θ such that M_θ is non-degenerate(i.e., $\prod_{i=1}^{N-1}\sin\theta_i \neq 0$),

$$H_{2\beta}^r(\theta) = \frac{1}{2^{N/(2\beta)}\theta_1^N}\Big(\prod_{i=1}^{N-1}\sin^{N-i}\theta_i\Big)H_{2\beta}.$$

Therefore, as $u \to \infty$,

$$\mathbb{P}\Big\{\sup_{x\in T} X(x) \geq u\Big\} \sim u^{N/\beta}\Psi(u)2^{-N/(2\beta)}H_{2\beta}\int_D \theta_1^{-N}\Big(\prod_{i=1}^{N-1}\sin^{N-i}\theta_i\Big)d\theta. \tag{2.10}$$

Remark 2.6 Comparing the excursion probabilities in (2.10) for the standardized SFBM X and in Theorem 2.4 for the isotropic Gaussian field W^β, which is defined in Estrade and Istas (2010), we see that the constant in (2.10) is more complicated.

3 Smooth Isotropic Gaussian Fields on Sphere

In this section we study the excursion probability of smooth isotropic Gaussian fields on sphere. Related to the results in this section, we mention that Cheng and Schwartzman (2014) have determined the height distribution and overshoot distribution of local maxima of smooth isotropic Gaussian random fields on sphere.

3.1 Preliminaries

Given $\lambda > 0$ and an integer $n \geq 0$, the *ultraspherical polynomial* (or *Gegenbauer polynomial*) of degree n, denoted by $P_n^\lambda(t)$, is defined by the expansion

$$(1 - 2rt + r^2)^{-\lambda} = \sum_{n=0}^{\infty} r^n P_n^\lambda(t), \quad t \in [-1, 1].$$

For $\lambda = 0$, we follow Schoenberg (1942) and define $P_n^0(t) = \cos(n \arccos t) = T_n(t)$, where T_n $(n \geq 0)$ are *Chebyshev polynomials of the first kind* defined by the expansion

$$\frac{1 - rt}{1 - 2rt + r^2} = \sum_{n=0}^{\infty} r^n T_n(t), \quad t \in [-1, 1].$$

For reference later on, we recall the following formulae on P_n^λ.

(i). For all $n \geq 0$, $P_n^0(1) = 1$, and if $\lambda > 0$ [cf. Szegö (1975, p. 80)],

$$P_n^\lambda(1) = \binom{n + 2\lambda - 1}{n}. \tag{3.1}$$

(ii). For all $n \geq 0$,

$$\frac{d}{dt} P_n^0(t) = n P_{n-1}^1(t), \tag{3.2}$$

and if $\lambda > 0$ [cf. Szegö (1975, p. 81)],

$$\frac{d}{dt} P_n^\lambda(t) = 2\lambda P_{n-1}^{\lambda+1}(t). \tag{3.3}$$

The following theorem by Schoenberg (1942) characterizes the covariance function of an isotropic Gaussian field on sphere [see also Gneiting (2013)].

Theorem 3.1 *Let $N \geq 1$, then a continuous function $C(\cdot, \cdot) : \mathbb{S}^N \times \mathbb{S}^N \to \mathbb{R}$ is the covariance of an isotropic Gaussian field on $\mathbb{S}^N$ if and only if it has the form*

$$C(x, y) = \sum_{n=0}^{\infty} a_n P_n^\lambda(\langle x, y \rangle), \quad x, y \in \mathbb{S}^N,$$

where $\lambda = (N - 1)/2$, $a_n \geq 0$ and $\sum_{n=0}^{\infty} a_n P_n^\lambda(1) < \infty$.

Remark 3.2 Note that for the case of $N = 1$ and $\lambda = 0$, $\sum_{n=0}^{\infty} a_n P_n^0(1) < \infty$ is equivalent to $\sum_{n=0}^{\infty} a_n < \infty$; while for $N \geq 2$ and $\lambda = (N-1)/2$, (3.1)implies that $\sum_{n=0}^{\infty} a_n P_n^\lambda(1) < \infty$ is equivalent to $\sum_{n=0}^{\infty} n^{N-2} a_n < \infty$.

When $N = 2$ and $\lambda = 1/2$, P_n^λ $(n \geq 0)$ become the *Legendre polynomials*. For more results on isotropic Gaussian fields on $\mathbb{S}^2$, we refer to Marinucci and Peccati (2011). Regularity and smoothness properties of Gaussian field $\{X(x),\, x \in \mathbb{S}^2\}$ have recently been obtained by Lang and Schwab (2013) in terms of the corresponding angular power spectrum.

The following statement (**A**1) is a smoothness condition for Gaussian fields on sphere. In Lemma 3.3 below, we show that it implies $X(\cdot) \in C^2(\mathbb{S}^N)$ a.s.

(**A**1). The covariance $C(\cdot,\cdot)$ of $\{X(x) : x \in \mathbb{S}^N\}$ satisfies

$$C(x,y) = \sum_{n=0}^{\infty} a_n P_n^\lambda(\langle x, y\rangle), \quad x, y \in \mathbb{S}^N,$$

where $\lambda = (N-1)/2$, $a_n \geq 0$ and, $\sum_{n=1}^{\infty} n^{N+8} a_n < \infty$ if $N \geq 2$, $\sum_{n=1}^{\infty} n^{10} a_n < \infty$ if $N = 1$.

Lemma 3.3 *Let $\{X(x) : x \in \mathbb{S}^N\}$ be an isotropic Gaussian field such that* (**A**1) *is fulfilled. Then $X(\cdot) \in C^2(\mathbb{S}^N)$ a.s.*

Proof We first consider $N \geq 2$. By Theorem 3.1, each $P_n^\lambda(\langle t, s\rangle)$ is the covariance of an isotropic Gaussian field on $\mathbb{S}^N$ and hence the Cauchy-Schwarz inequality implies

$$|P_n^\lambda(\langle x, y\rangle)| \leq P_n^\lambda(\langle x, x\rangle) = P_n^\lambda(1), \quad \forall x, y \in \mathbb{S}^N. \tag{3.4}$$

Combining (**A**1) with (3.1), (3.3) and (3.4), together with the fact $P_0^\lambda(t) \equiv 1$, we obtain that there exist positive constants M_1 and M_2 such that

$$\sup_{t\in[-1,1]} \sum_{n=0}^{\infty} a_n \Big| \Big(\frac{d^5}{dt^5} P_n^\lambda(t) \Big) \Big| \leq M_1 \sum_{n=5}^{\infty} a_n P_{n-5}^{\lambda+5}(1) \leq M_2 \sum_{n=1}^{\infty} n^{N+8} a_n < \infty.$$

This shows that $C(\cdot,\cdot) \in C^5(\mathbb{S}^N \times \mathbb{S}^N)$. The proof for $N = 1$ is similar once we apply both (3.2) and (3.3). Therefore, by arguments via charts [cf. Auffinger (2011)] and the results in Potthoff (2010) (though the results therein are for $X(\cdot) \in C^1$, they can be extended easily to the case of higher-order smoothness), we conclude that $X(\cdot) \in C^2(\mathbb{S}^N)$ a.s. □

By Schoenberg (1942) or Gneiting (2012), $C(\cdot,\cdot)$ is a covariance function on $\mathbb{S}^N$ for every $N \geq 1$ if and only if it has the form

$$C(x,y) = \sum_{n=0}^{\infty} b_n \langle x, y\rangle^n, \quad x, y \in \mathbb{S}^N,$$

where $b_n \geq 0$ and $\sum_{n=0}^{\infty} b_n < \infty$. Then similarly to (**A**1), we may state the smoothness condition (**A**1′) below for this special class of Gaussian fields on sphere.

(**A**1′). The covariance $C(\cdot,\cdot)$ of $\{X(x) : x \in \mathbb{S}^N\}$ satisfies

$$C(x,y) = \sum_{n=0}^{\infty} b_n \langle x,y\rangle^n, \quad x,y \in \mathbb{S}^N,$$

where $b_n \geq 0$ and $\sum_{n=0}^{\infty} n^5 b_n < \infty$.

We obtain below an analogue of Lemma 3.3. Since the proof is similar, it is omitted.

Lemma 3.4 *Let $\{X(x) : x \in \mathbb{S}^N\}$ be an isotropic Gaussian field such that (**A**1′) is fulfilled. Then $X(\cdot) \in C^2(\mathbb{S}^N)$ a.s.*

3.2 Excursion Probability

Let $\chi(A_u(X,\mathbb{S}^N))$ be the Euler characteristic of excursion set $A_u(X,\mathbb{S}^N) = \{x \in \mathbb{S}^N : X(x) \geq u\}$ [cf. Adler and Taylor (2007)]. Denote by $H_j(x)$ the Hermite polynomial of order j, i.e.,

$$H_j(x) = (-1)^j e^{x^2/2} \frac{d^j}{dx^j}\big(e^{-x^2/2}\big).$$

Let $\omega_j = \text{Area}(\mathbb{S}^j)$, the spherical area of the j-dimensional unit sphere $\mathbb{S}^j$, i.e.,

$$\omega_j = \frac{2\pi^{(j+1)/2}}{\Gamma\big(\frac{j+1}{2}\big)}.$$

Before stating our results, we need another regularity condition for the Gaussian field.

(**A**2). For each $x \in \mathbb{S}^N$, the joint distribution of $(X(x), \nabla X(x), \nabla^2 X(x))$ is non-degenerate.

Lemma 3.5 *Let $\{X(x) : x \in \mathbb{S}^N\}$ be a centered, unit-variance, isotropic Gaussian field satisfying (**A**1) and (**A**2). Then*

$$\mathbb{E}\{\chi(A_u(X,\mathbb{S}^N))\} = \sum_{j=0}^{N} (C')^{j/2} \mathcal{L}_j(\mathbb{S}^N) \rho_j(u),$$

where the constant C' is defined as

$$C' = \begin{cases} (N-1)\sum_{n=1}^{\infty} \binom{n+N-1}{N} a_n & \text{if } N \geq 2, \\ \sum_{n=1}^{\infty} n^2 a_n & \text{if } N = 1, \end{cases} \tag{3.5}$$

and where $\rho_0(u) = (2\pi)^{-1/2} \int_u^{\infty} e^{-x^2/2} dx$, $\rho_j(u) = (2\pi)^{-(j+1)/2} H_{j-1}(u) e^{-u^2/2}$ for $j \geq 1$ and, for $j = 0, 1, \ldots, N$,

$$\mathcal{L}_j(\mathbb{S}^N) = \begin{cases} 2\binom{N}{j} \frac{\omega_N}{\omega_{N-j}} & \text{if } N-j \text{ is even}, \\ 0 & \text{otherwise} \end{cases} \tag{3.6}$$

are the Lipschitz-Killing curvatures of $\mathbb{S}^N$.

Remark 3.6 In Lemma 3.5, if condition (**A**1) is replaced by (**A**1′), then it can be seen from the proof below that the result still holds with C' being replaced by

$$C' = \sum_{n=1}^{\infty} n b_n. \tag{3.7}$$

Proof of Lemma 3.5 By Theorem 12.4.1 in Adler and Taylor (2007), we only need to show that the Lipschitz-Killing curvatures induced by X on $\mathbb{S}^N$ are $\mathcal{L}_j(X, \mathbb{S}^N) = (C')^{j/2}\mathcal{L}_j(\mathbb{S}^N)$ for $j = 0, 1, \ldots, N$.

The Riemannian structure induced by X on $\mathbb{S}^N$ is defined as [cf. Adler and Taylor (2007, p. 305)]

$$g_{x_0}^{X,\mathbb{S}^N}(\xi_{x_0}, \sigma_{x_0}) := \mathbb{E}\{(\xi_{x_0} X) \cdot (\sigma_{x_0} X)\} = \xi_{x_0}\sigma_{x_0} C(x, y)|_{x=y=x_0}, \quad \forall x_0 \in \mathbb{S}^N,$$

where $\xi_{x_0}, \sigma_{x_0} \in T_{x_0}\mathbb{S}^N$, the tangent space of $\mathbb{S}^N$ at x_0. We may choose two smooth curves on $\mathbb{S}^N$, say $\gamma(t)$, $\tau(s)$, $t, s \in [0, 1]$, such that $\gamma(0) = \tau(0) = x_0$ and $\gamma'(0) = \xi_{x_0}, \tau'(0) = \sigma_{x_0}$. We first consider $N \geq 2$, then

$$\begin{aligned}
\xi_{x_0}\sigma_{x_0} C(x, y)|_{x=y=x_0} &= \frac{\partial}{\partial t}\frac{\partial}{\partial s} C(\gamma(t), \tau(s))|_{t=s=0} \\
&= \frac{\partial}{\partial t}\frac{\partial}{\partial s} \sum_{n=0}^{\infty} a_n P_n^{\lambda}(\langle \gamma(t), \tau(s) \rangle)|_{t=s=0} \\
&= \frac{\partial}{\partial t} \sum_{n=1}^{\infty} a_n (N-1) P_{n-1}^{\lambda+1}(\langle \gamma(t), x_0 \rangle)\langle \gamma(t), \sigma_{x_0} \rangle|_{t=0} \\
&= \sum_{n=2}^{\infty} a_n (N-1)(N+2) P_{n-2}^{\lambda+2}(\langle x_0, x_0 \rangle)\langle \xi_{x_0}, x_0 \rangle\langle x_0, \sigma_{x_0} \rangle \\
&\quad + \sum_{n=1}^{\infty} a_n (N-1) P_{n-1}^{\lambda+1}(\langle x_0, x_0 \rangle)\langle \xi_{x_0}, \sigma_{x_0} \rangle \\
&= \left(\sum_{n=1}^{\infty} a_n (N-1) P_{n-1}^{\lambda+1}(1) \right) \langle \xi_{x_0}, \sigma_{x_0} \rangle = C' \langle \xi_{x_0}, \sigma_{x_0} \rangle,
\end{aligned}$$

where the third and fourth equalities follow from (3.3), while the fifth equality is due to the facts $\langle x_0, x_0 \rangle = 1$ and $\langle \xi_{x_0}, x_0 \rangle = \langle \sigma_{x_0}, x_0 \rangle = 0$, since the vector x_0 is always orthogonal to its tangent space. The case $N = 1$ can be proved similarly once we apply (3.2) instead of (3.3).

Hence the induced metric is

$$g_{x_0}^{X,\mathbb{S}^N}(\xi_{x_0}, \sigma_{x_0}) = C' \langle \xi_{x_0}, \sigma_{x_0} \rangle, \quad \forall x_0 \in \mathbb{S}^N.$$

By the definition of Lipschitz-Killing curvatures, one has

$$\mathcal{L}_j(X, \mathbb{S}^N) = (C')^{j/2} \mathcal{L}_j(\mathbb{S}^N),$$

where $\mathcal{L}_j(\mathbb{S}^N)$ are the original Lipschitz-Killing curvatures of $\mathbb{S}^N$ given by (3.6). We have finished the proof. □

Applying Lemma 3.5 and Theorem 14.3.3 in Adler and Taylor (2007), we obtain immediately the following approximation for the excursion probability.

Theorem 3.7 *Suppose the conditions in Lemma 3.5 hold. Then, under the notation therein, there exists a constant $\alpha_0 > 0$ such that as $u \to \infty$,*

$$\mathbb{P}\Big\{ \sup_{x\in\mathbb{S}^N} X(x) \geq u \Big\} = \sum_{j=0}^{N} (C')^{j/2} \mathcal{L}_j(\mathbb{S}^N) \rho_j(u) + o(e^{-\alpha_0 u^2 - u^2/2}). \tag{3.8}$$

The following are some remarks.

Remark 3.8

- Under the conditions in Theorem 3.7, the covariance function C satisfies (2.1) with $\alpha = 2$. Also note that when $\alpha = 2$, Pickands' constant $H_2 = \pi^{-N/2}$. Then one can check that the approximation in Theorem 2.4 only provides the leading term of the approximation in Theorem 3.7. This also affects the errors in two approximations: the error in the former one is only $o(u^{N-1}e^{-u^2/2})$, while the error in the latter one is $o(e^{-\alpha_0 u^2 - u^2/2})$.

- By applying the tube method, Sun (1993) gave a two-term approximation formula for the excursion probability of a class of differentiable Gaussian random field $\{X(x), x \in I\}$, where $I \subset \mathbb{R}^N$ is a bounded convex set. Her results can be applied to provide a two-term approximation for the excursion probability in (3.8) for some special cases. See Park and Sun (1998, p. 73).

- Recently Marinucci and Vadlamani (2013) have computed the Lipschitz-Killing curvatures of excursion set and derived a very precise approximation for the excursion probability of a class of nonlinear functionals of a smooth Gaussian random field on $\mathbb{S}^2$. In the linear case (i.e., $q = 1$) Theorem 21 of Marinucci and Vadlamani (2013) is a special case of (3.8) with $N = 2$.

If the sphere $\mathbb{S}^N$ is replaced by a more general subset $T \subset \mathbb{S}^N$, by simply revising Lemma 3.5 and applying Theorem 14.3.3 in Adler and Taylor (2007) again, we obtain the following corollary.

Corollary 3.9 *Suppose the conditions in Lemma 3.5 hold. Let $T \subset \mathbb{S}^N$ be a k-dimensional, locally convex, regular stratified manifold [cf. Adler and Taylor (2007, p. 198)], then there*

exists $\alpha_0 > 0$ *such that as* $u \to \infty$,

$$\mathbb{P}\Big\{ \sup_{x\in T} X(x) \geq u \Big\} = \sum_{j=0}^{k} (C')^{j/2} \mathcal{L}_j(T) \rho_j(u) + o(e^{-\alpha_0 u^2 - u^2/2}), \tag{3.9}$$

where $\mathcal{L}_j(T)$ *are the Lipschitz-Killing curvatures of* T *[cf. Adler and Taylor (2007, p. 175)],* C' *and* $\rho_j(u)$ *are as in Lemma 3.5.*

The parameter set $T \subset \mathbb{S}^N$ in Corollary 3.9 is assumed to be nice enough. Roughly speaking, it looks like convex and can be decomposed into several smooth manifolds, see Adler and Taylor (2007) for rigorous definition. Also, the j-th Lipschitz-Killing curvature $\mathcal{L}_j(T)$ is like the measure of the j-dimensional boundary of T. One may use Steiner's formula [Adler and Taylor (2007, p. 142)] to compute the Lipschitz-Killing curvatures of T exactly. In particular, if T is a semisphere of dimension one, then $\mathcal{L}_0(T) = 1$ and $\mathcal{L}_1(T) = \pi$. If T is a semisphere of dimension two, then $\mathcal{L}_0(T) = 1$, $\mathcal{L}_1(T) = \pi$ and $\mathcal{L}_2(T) = 2\pi$. More generally, if T is a k-dimensional, locally convex, regular stratified manifold, then $\mathcal{L}_0(T)$ is the Euler characteristic, $\mathcal{L}_k(T)$ is the volume and $\mathcal{L}_{k-1}(T)$ is half of the surface area. For the other $\mathcal{L}_j(T)$, $1 \leq j \leq k-2$, we can apply Steiner's formula to find their values.

Lastly, to further illustrate the main results of this paper, we give more examples on approximating the excursion probability of Gaussian fields on spheres, including both smooth and non-smooth cases.

Example 3.10 The canonical Gaussian field on $\mathbb{S}^N$, denoted by X, has covariance function given by $C(x, y) = \langle x, y\rangle$ [cf. Adler and Taylor (2007)]. Since $C(x, y) = \cos d(x, y)$, it satisfies

$$C(x, y) = 1 - \frac{1}{2} d^2(x, y)(1 + o(1)), \quad \text{as } d(x, y) \to 0.$$

Applying Theorem 2.4 with $T = \mathbb{S}^N$, $c = 1/2$ and $\alpha = 2$, we obtain an approximation to the excursion probability:

$$\mathbb{P}\Big\{ \sup_{x\in\mathbb{S}^N} X(x) \geq u \Big\} \sim 2^{-N/2} \text{Area}(\mathbb{S}^N) H_2 u^N \Psi(u) = (2\pi)^{-(N+1)/2} \omega_N u^{N-1} e^{-u^2/2}.$$

However, by applying Theorem 3.7 with $C' = 1$, we get a more precise approximation:

$$\mathbb{P}\Big\{ \sup_{x\in\mathbb{S}^N} X(x) \geq u \Big\} = \sum_{j=0}^{N} \mathcal{L}_j(\mathbb{S}^N) \rho_j(u) + o(e^{-\alpha_0 u^2 - u^2/2}).$$

Example 3.11 Consider the Hamiltonian of the pure p-spin model on $\mathbb{S}^{N-1}$

$$H_{N,p}(x) = \frac{1}{N^{(p-1)/2}} \sum_{i_1,\ldots,i_p=1}^{N} J_{i_1,\ldots,i_p} x_{i_1} \cdots x_{i_p}, \quad \forall x = (x_1, \cdots, x_N) \in \mathbb{S}^{N-1},$$

where $J_{i_1,\ldots,i_p}$ are independent standard Gaussian random variables. Then

$$\mathbb{E}\{H_{N,p}(x)H_{N,p}(y)\} = \frac{1}{N^{p-1}}\langle x, y\rangle^p.$$

Let

$$X(x) = \sum_{p=2}^{\infty} b_p H_{N,p}(x),$$

where $(b_p)_{p\geq 2}$ is a sequence of positive numbers such that $\sum_{p=2}^{\infty} 2^p b_p < \infty$, and $H_{N,p}$ and $H_{N,p'}$ are independent for any $p \neq p'$. Then X is a smooth Gaussian random field on $\mathbb{S}^{N-1}$ with covariance

$$C(x, y) = \sum_{p=2}^{\infty} \frac{b_p^2}{N^{p-1}}\langle x, y\rangle^p,$$

and we can apply Theorem 3.7 or Corollary 3.9 to approximate the excursion probability.

Example 3.12 Consider the Gaussian field $\{X(x) : x \in \mathbb{S}^N\}$ with covariance structure $C(x, y) = 1 - \frac{2}{\pi}d(x, y)$, which can be verified to be a valid covariance on sphere [cf. Zuo (2003, Remark 3.3)]. Since $d(x, y) = \arccos\langle x, y\rangle$, we can write

$$\begin{aligned} C(x, y) = 1 - \frac{2}{\pi}\arccos\langle x, y\rangle &= \sum_{n=0}^{\infty} \frac{(2n)!}{4^n(n!)^2(2n+1)}\langle x, y\rangle^{2n+1} \\ &:= \sum_{n=0}^{\infty} b_n\langle x, y\rangle^n, \end{aligned} \tag{3.10}$$

it is easy to check that $\sum_{n=0}^{\infty} nb^n = \infty$, $(\mathbf{A}1')$ is not satisfied and hence Theorem 3.7 is not applicable. Instead, we may use Theorem 2.2 to get the approximation to excursion probability. This result allows one to construct confidence regions for the true projection median defined in Zuo (2003, Section 3) without using the bootstrapping techniques.

Acknowledgment. The authors thank Professor Enkelejd Hashorva, Dr. Lanpeng Ji for stimulating discussions and Dr. Xiaohui Liu for pointing out the connection between Gaussian random fields on sphere and projection depth functions in Zuo (2003).

References

[1] Adler, R. J. (2010). *The Geometry of Random Fields*. (SIAM Classics ed.). Philadelphia: SIAM.

[2] Adler, R. J. and Taylor, J. E. (2007), *Random Fields and Geometry*. Springer, New York.

[3] Adler, R. J., Taylor, J. E. and Worsley, K. J. (2012),*Applications of Random Fields and Geometry: Foundations and Case Studies*. In preparation.

[4] Andreev, R. and Lang, A. (2013), Kolmogorov-Chentsov theorem and differentiability of random fields on manifolds. Available at *http://arxiv.org/abs/1307.4886*.

[5] Auffinger, A. (2011), *Random Matrices, Complexity of Spin Glasses and Heavy Tailed Processes*. Ph.D. Thesis, New York University.

[6] Cabella, P. and Marinucci, D. (2009), Statistical challenges in the analysis of Cosmic Microwave Background radiation. *Ann. Appl. Stat.* **3**, 61–95.

[7] Chan, H. P. and Lai, T. L. (2006), Maxima of asymptotically Gaussian random fields and moderate deviation approximations to boundary crossing probabilities of sums of random variables with multidimensional indices. *Ann. Probab.* **34**, 80–121.

[8] Cheng, D. and Schwartzman, A. (2014), Distribution of the height of local maxima of Gaussian random fields. *Submitted*. Available at *http://arxiv.org/abs/1307.5863*

[9] Du, J., Ma, C. and Li, Y. (2013), Isotropic variogram matrix functions on spheres. *Math. Geosci.* **45**, 341–357.

[10] Estrade, A. and Istas, J. (2010), Ball throwing on spheres. *Bernoulli* **16**, 953–970.

[11] Gangolli, R. (1967), Positive definitive kernels on homogeneous spaces and certain stochastic processes related to Lévy Brownian motion of several parameters. *Ann. Inst. H. Poincaré Sect. B* **3**, 121–226.

[12] Genovese, C. R., Miller, C. J., Nichol, R C., Arjunwadkar, M. and Wasserman, L. (2004), Nonparametric inference for the cosmic microwave background. *Statist. Sci.* **19**, 308–321.

[13] Gneiting, T. (2013), Strictly and non-strictly positive definite functions on spheres. *Bernoulli* **19**, 1327–1349.

[14] Hansen, L. V., Thorarinsdottir, T. L. and Gneiting, T. and Richards, D. (2012). Lévy particles: Modelling and simulating star-shaped random sets. Available at *http://sites.stat.psu.edu/˜richards/papers/aap4a.pdf.*

[15] Hitczenko, M. and Stein, M. L. (2012), Some theory for anisotropic processes on the sphere. *Stat. Methodol.* **9**, 211–227.

[16] Huang. C., Zhang, H. and Robeson S. M. (2011). On the validity of commonly used covariance and variogram functions on the sphere. *Math. Geosci.* **43**, 721–733.

[17] Istas, J. (2005), Spherical and hyperbolic fractional Brownian motion. *Electron. Comm. Probab.* **10**, 254–262.

[18] Istas, J. (2006), Karhunen-Loève expansion of spherical fractional Brownian motions. *Statist. Probab. Lett.* **76**, 1578–1583.

[19] Jones, R. H. (1963), Stochastic processes on a sphere. *Ann. Math. Statist.* **34**, 213–218.

[20] Jun, M. (2011), Non-stationary cross-covariance models for multivariate processes on a globe. *Scand. J. Statist.* **38**, 726–747.

[21] Jun, M. and Stein, M. L. (2007), An approach to producing space-time covariance functions on spheres. *Technometrics* **49**, 468–479.

[22] Jun, M. and Stein, M. L. (2008), Nonstationary covariance models for global data. *Ann. Appl. Statist.* **2**, 1271–1289.

[23] Lang, A. and Schwab, C. (2013), Isotropic Gaussian random fields on the sphere: regularity, fast simulation, and stochastic partial differential equations. Available at *http://arxiv.org/abs/1305.1170*.

[24] Ma, C. (2012), Stationary and isotropic vector random fields on spheres. *Math. Geosci.* **44**, 765–778.

[25] Marinucci, D. and Peccati, G. (2011), *Random Fields on the Sphere: Representation, Limit Theorems and Cosmological Applications.* Cambridge University Press, Cambridge, UK.

[26] Marinucci, D. and Vadlamani, S. (2013), High-frequency asymptotics for Lipschitz-Killing curvatures of excursion sets on the sphere. Available at *http://arxiv.org/pdf/1303.2456v1.pdf*.

[27] Mikhaleva, T. L. and Piterbarg, V. I. (1997), On the distribution of the maximum of a Gaussian field with constant variance on a smooth manifold. *Theory Probab. Appl.* **41**, 367–379.

[28] Obukhov, A. M. (1947), Statistically homogeneous fields on a sphere. *Uspekhi math. Nauk* **2**, 196–198.

[29] Oh, H.-S. and Li, T.-H. (2004), Estimation of global temperature fields from scattered observations by a spherical-wavelet-based spatially adaptive method. *J. Roy. Statist. Soc. Ser. B* **66**, 221–238.

[30] Park, M. G. and Sun, J. (1998), Tests in projection pursuit regression. *J. Statist. Plann. Inference* **75**, 65–90.

[31] Piterbarg, V. I. (1996), *Asymptotic Methods in the Theory of Gaussian Processes and Fields.* Translations of Mathematical Monographs, Vol. 148, American Mathematical Society, Providence, RI.

[32] Pickands, J. (1969), Upcrossing probabilities for stationary Gaussian processes. *Trans. Amer. Math. Soc.* **145**, 51–75.

[33] Potthoff, J. (2010), Sample properties of random fields III: differentiability. *Commun. Stoch. Anal.* **4**, 335–353.

[34] Qualls, C. and Watanabe, H. (1973), Asymptotic properties of Gaussian random Fields. *Trans. Amer. Math. Soc.* **177**, 155–171.

[35] Schoenberg, I. J. (1942), Positive definite functions on spheres. *Duke Math. J.* **9**, 96–108.

[36] Soubeyrand, S., Enjalbert, J. and Sache, I. (2008), Accounting for roughness of circular processes: Using Gaussian random processes to model the anisotropic spread of airborne plant disease. *Theoret. Population Bio.* **73**, 92–103.

[37] Stein, M. L. (2007), Spatial variation of total column Ozone on a global scale. *Ann. Appl. Statist.* **1**, 191–210.

[38] Sun, J. (2001), Multiple comparisons for a large number of parameters. *Biom. J.* **43**, 627–643.

[39] Sun, J. (1993), Tail probabilities of the maxima of Gaussian random fields. *Ann. Probab.* **21**, 34–71.

[40] Sun, J. (1991), Significance levels in exploratory projection pursuit. *Biometrika* **78**, 759–769.

[41] Szegö, G. (1975), *Othogonal Polynomials*. American Mathematical Society, Providence, RI.

[42] Tebaldi, C. and Sansó, B. (2009), Joint projections of temperature and precipitation change from multiple climate models: a hierarchical Bayesian approach. *J. Roy. Statist. Soc. Series A* **172**, 83–106.

[43] Yadrenko, A. M. (1983). *Spectral Theory of Random Fields.* Optimization Software, New York.

[44] Yaglom, A. M. (1961), Second-order homogeneous random fields. *Proc. Fourth Berkeley Symp. on Math. Statist. and Probab.*, Vol. 2, pp. 593–622. University of Calif. Press.

[45] Zuo, Y. (2003), Projection-based depth functions and associated medians. *Ann. Statist.* **31**, 1460–1490.